## **BRIEF REPORT**



# A whole food, plant-based diet reduces amino acid levels in patients with metastatic breast cancer

TashJaé Q. Scales<sup>1,2</sup>, Bradley Smith<sup>2</sup>, Lisa M. Blanchard<sup>3</sup>, Nellie Wixom<sup>4</sup>, Emily T. Tuttle<sup>1,2</sup>, Brian J. Altman<sup>1,2</sup>, Luke J. Peppone<sup>2,5</sup>, Joshua Munger<sup>2,6</sup>, Thomas M. Campbell<sup>3</sup>, Erin K. Campbell<sup>7</sup> and Isaac S. Harris<sup>1,2\*</sup>

## Abstract

**Background** Amino acids are critical to tumor survival. Tumors can acquire amino acids from the surrounding microenvironment, including the serum. Limiting dietary amino acids is suggested to influence their serum levels. Further, a plant-based diet is reported to contain fewer amino acids than an animal-based diet. The extent to which a plantbased diet lowers the serum levels of amino acids in patients with cancer is unclear.

**Methods** Patients with metastatic breast cancer (n = 17) were enrolled in a clinical trial with an ad libitum whole food, plant-based diet for 8 weeks without calorie or portion restriction. Dietary changes by participants were monitored using a three-day food record. Serum was collected from participants at baseline and 8 weeks. Food records and serum were analyzed for metabolic changes.

**Results** We found that a whole food, plant-based diet resulted in a lower intake of calories, fat, and amino acids and higher levels of fiber. Additionally, body weight, serum insulin, and IGF were reduced in participants. The diet contained lower levels of essential and non-essential amino acids, except for arginine (glutamine and asparagine were not measured). Importantly, the lowered dietary intake of amino acids translated to reduced serum levels of amino acids in participants (5/9 essential amino acids; 4/11 non-essential amino acids).

**Conclusions** These findings provide a tractable approach to limiting amino acid levels in persons with cancer. This data lays a foundation for studying the relationship between amino acids in patients and tumor progression. Further, a whole-food, plant-based diet has the potential to synergize with cancer therapies that exploit metabolic vulnerabilities.

**Trial registration** The clinical trial was registered with ClinicalTrials.gov identifier NCT03045289 on 2017–02-07. **Keywords** Whole food, Plant-based, Amino acids, Cancer, Diet, Fat, Carbohydrate, Protein, Fiber, Calories

\*Correspondence: Isaac S. Harris isaac\_harris@urmc.rochester.edu Full list of author information is available at the end of the article



© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/.

## Background

To grow and survive, cancer cells face increased metabolic demands, including a heightened need for amino acids [1, 2]. Tumors obtain amino acids through either intracellular production or extracellular uptake. The expression and activity of key enzymes associated with amino acid production are elevated in tumors compared with non-tumorigenic tissue [3, 4]. While numerous efforts have explored blocking these enzymes for a therapeutic benefit to cancer patients [5-9], they have met minimal clinical success. A potential explanation for the lack of efficacy of these drugs in clinical settings is the compensatory uptake of circulating amino acids by tumors. An alternative strategy is to lower the levels of amino acids in circulation [10–13], thus limiting the tumor's ability to uptake amino acids. Indeed, decreasing the dietary intake of either total protein [14–17] or specific amino acids [18–21] significantly perturbs tumor growth and progression in pre-clinical models of cancer. Although consumption of a healthy diet correlates with lower cancer risk [22], there is a lack of knowledge of how diet impacts amino acid levels in patients with cancer. Even more, it is unclear whether adopting a diet with reduced amino acids is feasible for patients with cancer.

A plant-based diet lower in processed foods has several potential anti-cancer benefits compared to a Western diet (i.e., an animal-based diet higher in processed foods). Higher fiber intake, commonly associated with a plantbased diet, improves a tumor's response to immunotherapy [23]. A plant-based diet also contains less fats, which are directly associated with increased tumor incidence and progression [24-26]. Additionally, a low-fat plantbased diet is associated with lower energy intake than a low-carbohydrate animal-based diet [27], suggesting a plant-based diet could potentially limit energy sources available to tumors. Finally, a plant-based diet contains less protein than animal-based diets [28, 29], suggesting a potential approach to deprive tumors of amino acids. However, whether the implementation of a plant-based diet can lower amino acid levels in patients with cancer is largely unknown.

In this study, a whole food, plant-based diet was adhered to by patients with metastatic breast cancer. While this was an ad-libitum diet with no restriction on food consumption, we found that this diet lowered the intake of calories, fat, and total protein in participants. Further, we showed reduced serum insulin and IGF levels, suggesting an approach to limiting pro-tumorigenic signals. Finally, we found that participants who consumed a whole food, plant-based diet had lower levels of amino acids in their serum. Together, these findings suggest that adopting a whole food, plant-based diet is a tractable approach towards lowering circulating amino acid levels in patients with cancer and potentially depleting cancers of key metabolites required for their growth and survival.

### Methods

#### **Clinical study selection**

Demographic details, recruitment details, and cardiometabolic outcomes have been previously described [30]. The cohort for this study consisted of patients with metastatic breast cancer and varying status of estrogen-receptor positive, progesterone receptor positive, or HER2 positive. The patients selected were on a stable treatment for  $\geq 6$  weeks and had no planned upcoming changes to their cancer therapy. The patients were screened to meet the requirements needed to participate, such as exclusions for patients with excessive alcohol intake, smoking, use of certain medications such as blood thinners, or any alteration that could be a confounding factor, such as recent surgery or recent implementation of the vegan diet. The screening for the participants consisted of a group information session regarding the dietary intervention and an individual visit to review medical history and confirm eligibility for the study. If the participants did not complete the screening or meet eligibility requirements, they were excluded from the study. Participants were randomized into two cohorts with blocks of 4 for the clinical study. The participants on the ad libitum whole food plant-based diet (n=21) were given three meals and a side dish daily for a total of 8 weeks. Details regarding diet intervention have been previously described [31]. Briefly, participants followed a whole food, plant-based (WFPB) diet with unrestricted eating. This diet included a variety of plant-based foods such as fruits, vegetables, whole grains, beans, nuts, and seeds. Soy products were permitted, and participants could consume small quantities of added sugars. However, the diet strictly prohibited any animal-derived foods, cooking oils, and solid fats.

Four intervention subjects were not included in this analysis. One forgot to complete their final three-day food record, another was withdrawn from the study by investigators early in their participation due to the pandemic shutdown, two were deemed poorly compliant with the study diet, one fully non-compliant, by predefined criteria, and one only borderline compliant with high intake of animal-based foods. The control group (n=11) were asked to maintain their regular diet and were not included in this analysis. Participants attended weekly visits with study physicians and completed assessments at baseline, week 4, and week 8.

### Ad libtum whole food plant-based diet

Patients could consume their own foods in addition to, or in place of, the provided meals if their foods were

'on plan.' Participants were asked to consume as much as they wanted and eat as frequently as they wanted to feel satiated. Assessment of the patient's dietary intake consisted of a 3-day food record at baseline and again at 8 weeks and three unscheduled 24-h phone calls from a dietitian. The 3-day record consisted of two weekdays and one weekend day. The records were analyzed using the Nutrition Data System for Research (NDSR) version 2017 (Nutrition Coordinating Center, University of Minnesota, Minneapolis, MN). To calculate the percent difference in dietary components (i.e., consumption, calories, fat, carbohydrates, total protein, animal protein, vegetable protein, fiber, and amino acids), an average value was calculated based on the 3 daily food records at baseline. For the percent difference at baseline, the difference between the baseline value and the baseline average was divided by the baseline average and multiplied by 100%. For the percent difference at the 8-week follow-up, the difference between the 8-week value and the baseline average was divided by the baseline average and multiplied by 100%.

#### Serum analysis

Serum samples for this analysis were collected at baseline and week 8. Serum samples (20  $\mu$ l) were added to an extraction solution (1000  $\mu$ l; 80% methanol with internal standards), vortexed, and centrifuged at max speed for 5 min. The supernatant (900  $\mu$ l) was transferred to a new microcentrifuge tube and stored at -80 C. For metabolite preparation, 900  $\mu$ L of metabolite extracts were dried with N<sub>2</sub> gas and reconstituted in 90  $\mu$ L of 50% acetonitrile (A955, Fisher Scientific).

For LC-MS analysis, metabolite extracts were analyzed by high-resolution mass spectrometry with an Orbitrap Exploris 240 (Thermo) coupled to a Vanquish Flex liquid chromatography system (Thermo). 2 µL of samples were injected into a Waters XBridge XP BEH Amide column (150 mm length × 2.1 mm id, 2.5 µm particle size) maintained at 25C, with a Waters XBridge XP VanGuard BEH Amide (5 mm  $\times$  2.1 mm id, 2.5 µm particle size) guard column. Mobile phase A was 100% LC-MS grade H<sub>2</sub>O with 20 mM ammonium formate and 0.1% formic acid. Mobile phase B was 90% acetonitrile with 20 mM ammonium formate and 0.1% formic acid. The gradient was 0 min, 100% B; 2 min, 100% B; 3 min, 90% B; 5 min, 90% B; 6 min, 85% B; 7 min, 85% B; 8 min, 75% B; 9 min, 75% B; 10 min, 55% B; 12 min, 55% B; 13 min, 35%, 20 min, 35% B; 20.1 min, 35% B; 20.6 min, 100% B; 22.2 min, 100% all at a flow rate of 150  $\mu$ l min – 1, followed by 22.7 min, 100% B; 27.9 min, 100% B at a flow rate of 300  $\mu$ l min – 1, and finally 28 min, 100% B at flow rate of 150  $\mu$ l min – 1, for a total length of 28 min. The H-ESI source was operated in positive mode at spray voltage 3500 with the following parameters: sheath gas 35 au, aux gas 7 au, sweep

gas 0 au, ion transfer tube temperature 320 C, vaporizer temperature 275 C, mass range 70 to 800 m/z, full scan MS1 mass resolution of 120,000 FWHM, RF lens at 70%, and standard automatic gain control (AGC). LC–MS data were analyzed by El-Maven software [32], and compounds were identified by matching to LC–MS methodspecific retention time values of external standards. Samples were run in two batches. To calculate relative abundance, peak intensities for baseline and 8-week follow-up samples were divided by the average peak intensity at baseline within each batch.

#### Results

## A whole food, plant-based diet decreases the consumption of calories, fat, and protein in patients with metastatic breast cancer

A whole food, plant-based diet is reported to contain less protein than traditional animal-based diets (i.e., Western diets) [28, 29, 33]. Patients with metastatic breast cancer enrolled in clinical trial NCT03045289 were provided an ad-libitum whole food, plant-based diet for 8 weeks [30, 31]. Notably, participants were very adherent to the diet and reported positive outcomes regarding perceived cognitive function, emotional well-being, fatigue, and hunger [31]. Participants documented their diets using three-day food records and unannounced 24-h food recalls, and serum was collected at baseline and 8 weeks (Fig. 1a). Analysis of food records and recalls found that participants consumed more total mass (including liquids) but less calories and fat (Fig. 1b-1d). The consumption of carbohydrates by participants trended towards an increase but was non-significant (Fig. 1e). These findings demonstrate that the adoption of a whole food, plant-based diet lowers the intake of calories and fat in patients with metastatic breast cancer. Since both caloric and fat consumption are predictors of tumor growth and progression [24-26], this suggests that a whole food, plant-based diet can potentially benefit patients.

In pre-clinical models, plant-based diets are reported to lower the consumption of total protein (i.e., amino acids) [29]. In line with these findings, we found that participants adhering to a whole food, plant-based diet consumed significantly less protein (Fig. 1f). Further, there was a near-complete substitution of animal protein with vegetable protein (Fig. 1g-h). These findings demonstrate that the adoption of whole food, plantbased diet is a clear approach to lowering the consumption of amino acids in patients with cancer. Since cancers have an elevated demand for amino acids [1, 2], the consumption of a whole food, plant-based diet is potentially an opportunity to deprive cancers of these amino acids.



Fig. 1 A whole food, plant-based diet decreases the consumption of calories, fat, and protein in patients with metastatic breast cancer. A Patients with metastatic breast cancer adhered to a whole food plant-based diet for 8 weeks. Two 3-day food records and three unannounced 24-h recalls documented their dietary intake. Serum was collected at baseline and 8 weeks. B-I The percent difference between baseline and 8-week follow-up was measured in participants' (n=51; 17 participants each with 3 daily food records at baseline and 8-week follow-up) consumption of total mass (B), calories (C), fat (D), carbohydrates (E), total protein (F), animal protein (G), vegetable protein (H), and fiber (I) following the adoption of a whole food, plant-based diet. An unpaired t test was used to determine statistical significance. \*\* P value < 0.01, \*\*\*\* P value < 0.0001

Increased dietary fiber can support improved responses to tumor immunotherapy [23], and participants were found to consume increased amounts of total dietary fiber (Fig. 1i). This suggests that, along with decreasing the consumption of pro-tumorigenic metabolites (i.e., fat and protein), adopting a whole food, plant-based diet increases the consumption of metabolites (i.e., fiber) that support anti-cancer therapies.

## Consumption of a whole food, plant-based diet decreases serum levels of cholesterol, insulin, and IGF

A diet high in whole foods includes those lower in energy density, thus leading to lower calorie intake [27, 31, 34].

Additionally, the consumption of whole food results in increased calories lost in feces (i.e., lower metabolizable energy) [35], which could support weight loss. Further, increased consumption of vegetables, such as with the dietary approaches to stop hypertension (DASH) diet, is well-established to be beneficial towards improved metrics of cardiovascular health (i.e., lower levels of cholesterol, HDL, and LDL) [36, 37]. In line with the decreased caloric intake (Fig. 1c), metastatic breast cancer patients consuming a whole food, plant-based diet showed decreases in body mass index (BMI) and weight (Fig. 2a-b). While no differences in fasting serum triglycerides were observed (Fig. 2c), patients had lower levels

of serum cholesterol, HDL, and LDL (Fig. 2d-f). These results demonstrate that the adoption of a whole food, plant-based diet by patients with cancer supports weight loss and lowers prognostic markers of cardiovascular disease.

Numerous dietary factors [37–40], including protein intake [41–46], are reported to impact the abundance of growth hormones, such as insulin and insulin-like growth factor (IGF). These growth factors can signal molecular pathways to drive cancer growth and survival [47, 48]. Thus, limiting dietary protein would potentially lower the growth factors involved in glucose metabolism (i.e., insulin and IGF), impair cancer growth, and sensitize tumors with anti-cancer therapies, such as PI3K inhibitors [49]. Less is known whether lowering dietary proteins through a plantbased diet can lower levels of these growth factors. We found that participants consuming a whole-food, plant-based diet had lower levels of insulin and IGF (Fig. 2g-h). Further, these reductions in insulin and IGF occurred not only in the absence of a decrease in dietary carbohydrates but in a non-significant but trending increase in dietary carbohydrates (Fig. 1f). These results demonstrate that adopting a whole food, plantbased diet is a potential approach to lowering serum levels of tumor-promoting growth factors (i.e., insulin and IGF).



**Fig. 2** Consumption of a whole food, plant-based diet decreases serum levels of cholesterol, insulin, and IGF. **A-J** Body mass index (BMI) (**A**) and weight (**B**), and serum levels of triglycerides (**C**), total cholesterol (**D**), HDL (**E**), LDL (**F**), insulin (**G**), IGF (**H**), CA15-3 (**I**), and CA27.29 (**J**) were measured in patients at baseline and 8-week following the adoption of a whole food, plant-based diet. *n* = 17 for (A-H) and *n* = 16 for (**I-J**). A paired t test was used to determine statistical significance. ns = not significant, \* *P* value < 0.05, \*\*\* *P* value < 0.001, \*\*\*\* *P* value < 0.001

Limiting the dietary intake of total protein [14–17] or specific amino acids [18–21] has been shown to impede tumor growth in preclinical cancer models. While a change in tumor volume was not a primary endpoint of the 8-week clinical trial [31], we sought to examine if there were any potential changes in circulating tumor markers following the adoption of the diet. Serum markers CA15-3 and CA27.29, which correlate with breast tumor burden [50], showed non-significant but trending decreases in patients following the implementation of a whole food, plant-based diet (Fig. 2i-j). Future clinical trials with extended time frames (i.e., greater than 8 weeks) are required to better understand how a whole food, plant-based diet impacts tumor progression in patients.

## A whole food, plant-based diet reduces the levels of serum amino acids

Plant-based diets are reported to contain lower amounts of protein compared to animal-based diets [28]. Notably, equivalent reductions are described to occur across all amino acid species with a plant-based diet [29]. In line with these studies, we found that patients on a whole food, plant-based diet consumed fewer amino acids, which occurred to both essential and non-essential amino acids (Fig. 3a-c). However, a broad range of reductions in amino acid species was observed, some with minimal and non-significant reductions (i.e., arginine and aspartate) and others with dramatic reductions (i.e., methionine and proline). Further, due to the technical approaches for quantification of amino acids in foods, the levels of glutamine and asparagine in the diets were not determined. These findings show that the consumption of a whole food, plant-based diet by patients with metastatic breast cancer significantly lowers their intake of essential and non-essential amino acids.

Essential amino acid levels are suggested to be maintained through dietary intake. Thus, we examined whether lowered consumption of essential amino acids in the diet translated to lower levels in the serum of patients. Interestingly, while we found reductions, this was not the case for every essential amino acid, as only 5/9 were significantly lowered (i.e., isoleucine, leucine, lysine, phenylalanine, and threonine) (Fig. 3b). Certain amino acids (i.e., tryptophan and methionine) saw reductions but were not statistically significant. An outlier was valine, which had increased levels in the serum. These findings demonstrate that while essential amino acids are described to be solely maintained from dietary sources, reductions in the dietary intake of these amino acids do not predict reductions in the serum. Nonetheless, this shows that reducing essential amino acids in patients with cancer is clinically achievable and may contribute to slower tumor growth.

Unlike essential amino acids, the levels of non-essential amino acids are controlled by de novo synthesis and dietary intake. Thus, it was unclear if limiting non-essential acids would result in lower levels in patients. We found 4/11 non-essential amino acids (i.e., alanine, glutamate, proline, tyrosine) were lower in the serum of participants consuming a whole food, plant-based diet (Fig. 3c). Some of these reductions in non-essential amino acids could be due to reductions in essential amino acids; however, it was unclear why certain amino acids were reduced and others weren't (i.e., cystine, serine, glycine). Nonetheless, these results show that, along with reductions in serum levels of essential amino acids, the consumption of a whole food, plant-based diet leads to reductions in serum levels of non-essential amino acids in patients with cancer.

#### Discussion

Limiting the dietary intake of amino acids has been shown to slow tumor growth in pre-clinical models of cancer. We show that consuming a whole food, plantbased diet lowers the dietary intake and circulating levels of amino acids in patients with cancer. These studies provide a framework for clinically exploring the efficacy of lowering amino acids on slower tumor growth. Notably, significant efforts have been placed on limiting non-essential amino acids from the diet (i.e., serine and glycine) [51], but few studies have explored the possibility of limiting essential amino acids in patients, potentially due to the perception that limiting essential acids could be harmful. We find lowering both essential and nonessential amino acids in patients with cancer does not have a harmful impact; instead, participants reported a positive impact on several reported outcomes, including perceived cognitive function, emotional well-being, and

<sup>(</sup>See figure on next page.)

**Fig. 3** A whole food, plant-based diet reduces the levels of serum amino acids. **A-C** The percent difference in amino acid consumption between average and 8-week follow-up (**A**) and the relative abundance of essential (**B**) and non-essential (**C**) amino acids in the serum of patients at baseline and 8-week follow-up. An unpaired t test was used to determine the statistical significance of the percent difference in dietary consumption of amino acids (n = 51; 17 participants each with 3 daily food records at baseline and 8-week follow-up). A paired t test was used to determine the statistical significance of the relative abundance in serum levels of amino acids (n = 17). ns = not significant, \* *P* value < 0.05, \*\* *P* value < 0.01, \*\*\* *P* value < 0.001, \*\*\*\* *P* value < 0.001



fatigue [31]. However, one consideration is that, compared to preclinical studies [18–21], this diet lowered these amino acids rather than eliminated them. Further, reductions in dietary protein and total calories from this diet may lead to loss of muscle mass, and the lack of body composition measures limits the current study. Future clinical trials, with extended time frames, are required to examine if lowering these amino acids impacts tumor growth in patients.

Lowering circulating growth factors (i.e., insulin and IGF) is a therapeutic approach to improving anti-cancer drug efficacy [49], and dietary protein abundance can drive the production of these growth factors [44, 45]. We found that adopting a whole food, plant-based diet reduced the levels of insulin and IGF in participants. This occurred with a non-significant but trending increase in dietary carbohydrates. These findings suggest a clear approach to limiting these pro-tumor growth factors that do not require limiting the consumption of carbohydrates. Further, it suggests that combining this diet with anti-cancer therapies targeting these signaling pathways (i.e., PI3K inhibitors) would potentially act synergistically in slower tumor growth.

Essential amino acids cannot be synthesized and must be consumed from the diet. We found that a whole food, plant-based diet lowered the dietary intake of every essential amino acid. Surprisingly, this did not translate into a corresponding reduction in every essential amino acid in the serum of patients. Notably, levels of valine showed an increase in levels, which is counterintuitive. Further research, including a longer trial duration and larger sample size, is required to better understand how the levels of essential amino acids are regulated in the body and impacted by diet.

Following the implementation of a whole food, plantbased diet, we observed decreased serum amino acid levels in participants, including glutamate, isoleucine, and methionine. However, based on the degree of reductions and limited duration of the study, the extent to which the decreased serum amino acid levels would influence cancer progression was unclear. Previous studies have demonstrated that a glutamine restriction diet slows cancer progression [52]; notably, this diet lowered serum glutamate levels to a similar extent that was observed with a whole food, plant-based diet. Mice provided with a low isoleucine diet have a reduced incidence of tumor onset [53]. While no reductions in serum isoleucine levels were observed in mice on a low isoleucine diet, we found that patients on a whole food, plant-based diet showed reductions in serum isoleucine levels. Mice provided a methionine-restricted diet have lower serum methionine levels and slower tumor growth [20]. Here, the reduction in serum methionine in mice following a methionine-restricted diet was greater than the reduction in serum methionine observed in patients following a whole food, plant-based diet. However, the feasibility of a methionine-restricted diet for patients is unclear. Notably, clinical trials examining the implementation of a methionine-restricted have terminated due in part to slow accrual (NCT03186937, NCT03733119). Overall, in comparison to previous pre-clinical studies, it would suggest that the reductions in serum amino acid levels in patients following a whole food, plant-based diet would potentially slow cancer progression.

Numerous diets are suggested to deprive tumors of nutrients and improve outcomes in patients with cancer [54]. However, the probability of patients adopting these diets is more limited. We found that switching from an animal-based diet (i.e., Western diet) to a whole food, plantbased diet was feasible without any limitations to patients' quality of life. On the contrary, participants reported positive outcomes from the diet. A potential explanation is that this diet was provided ad libitum, without any limitation in food intake, and participants consumed more total food mass. Further, since this diet requires no special formulation (i.e., protein shakes), it is a highly tractable approach. Overall, this data suggests that the consumption of a whole food, plant-based diet is a clear method of lowering circulating amino acids in patients with cancer and potentially an approach to deprive tumors of nutrients required for their growth and survival.

#### Abbreviations

IGF	Insulin-like growth factor
HER2	Human epidermal growth factor receptor 2
NDSR	Nutrition data system for research
LC–MS	Liquid chromatography-mass spectrometry
DASH	Dietary approaches to stop hypertension
BMI	Body mass index
HDL	High-density lipoprotein
LDL	Low-density lipoprotein
PI3K	Phosphoinositide 3-kinase
CA15-3	Cancer antigen 15–3
CA27-29	Cancer antigen 27–29

#### Acknowledgements

We would like to thank the Metabolomics Resource and the Center for Advanced Research Technologies (CART) at URMC. The schematic in Figure 1A was created in BioRender. Harris, I. (2024) https://BioRender.com/p06w548.

#### Authors' contributions

T.Q.S. and I.S.H. initiated the study and conceived the project, designed experiments, interpreted results, and wrote the manuscript. B.S. and J.M. performed metabolite analysis. L.M.B., N.W., T.M.C., and E.K.C. conducted the clinical trial and sample acquisition. E.T.T., B.J.A., and L.J.P. supported project administration and provided expert comments.

#### Funding

This work was supported by Wilmot Cancer Institute pilot funding (I.S.H.), American Cancer Society grant RSG-23–971782-01-TBE, NIH grants R01CA269813 (I.S.H.) and R01CA269813-S2 (T.Q.S.), and the Highland Hospital Foundation (T.M.C. and E.K.C), which has received philanthropic donations from the T. Colin Campbell Center for Nutrition Studies, the Ladybug Foundation, and multiple individual donors.

#### Data availability

No datasets were generated or analysed during the current study.

#### Declarations

#### Ethics approval and consent to participate

This clinical study occurred at the University of Rochester Medical Center and complied with recognized ethical guidelines, including the U.S. Common Rule. The study protocol was approved by the University of Rochester Research Subject Review Board (ClinicalTrials.gov identifier: NCT03045289), and written informed consent was obtained from all participants.

#### **Consent for publication**

Not applicable.

#### **Competing interests**

The authors declare no competing interests.

#### Author details

<sup>1</sup>Department of Biomedical Genetics, University of Rochester Medical Center, Rochester, NY 14642, USA. <sup>2</sup>Wilmot Cancer Institute, University of Rochester Medical Center, Rochester, NY 14642, USA. <sup>3</sup>Department of Family Medicine, University of Rochester Medical Center, 601 Elmwood Ave, Rochester, NY 14642, USA. <sup>4</sup>Clinical Research Center, University of Rochester Medical Center, Rochester, NY, USA. <sup>5</sup>Department of Surgery, Cancer Control, University of Rochester Medical Center, Rochester, NY 14642, USA. <sup>6</sup>Department of Biochemistry and Biophysics, University of Rochester Medical Center, NY 14642, USA. <sup>7</sup>Department of Public Health Sciences, University of Rochester Medical Center, 601 Elmwood Ave, Rochester, NY 14642, USA.

#### Received: 21 October 2024 Accepted: 11 December 2024 Published online: 19 December 2024

#### References

- 1. DeBerardinis RJ, Chandel NS. Fundamentals of cancer metabolism. Sci Adv. 2016;2(5):e1600200.
- Martinez-Reyes I, Chandel NS. Cancer metabolism: looking forward. Nat Rev Cancer. 2021;21(10):669–80.
- Mendez-Lucas A, Lin W, Driscoll PC, Legrave N, Novellasdemunt L, Xie C, et al. Identifying strategies to target the metabolic flexibility of tumours. Nat Metab. 2020;2(4):335–50.
- Possemato R, Marks KM, Shaul YD, Pacold ME, Kim D, Birsoy K, et al. Functional genomics reveal that the serine synthesis pathway is essential in breast cancer. Nature. 2011;476:346–50.
- Gross MI, Demo SD, Dennison JB, Chen L, Chernov-Rogan T, Goyal B, et al. Antitumor activity of the glutaminase inhibitor CB-839 in triple-negative breast cancer. Mol Cancer Ther. 2014;13(4):890–901.
- Ngo B, Kim E, Osorio-Vasquez V, Doll S, Bustraan S, Liang RJ, et al. Limited Environmental Serine and Glycine Confer Brain Metastasis Sensitivity to PHGDH Inhibition. Cancer Discov. 2020;10(9):1352–73.
- Pacold ME, Brimacombe KR, Chan SH, Rohde JM, Lewis CA, Swier LJYM, et al. A PHGDH inhibitor reveals coordination of serine synthesis and onecarbon unit fate. Nat Chem Biol. 2016;12:452.
- Choi BH, Rawat V, Hogstrom J, Burns PA, Conger KO, Ozgurses ME, et al. Lineage-specific silencing of PSAT1 induces serine auxotrophy and sensitivity to dietary serine starvation in luminal breast tumors. Cell Rep. 2022;38(3): 110278.
- Lukey MJ, Cluntun AA, Katt WP, Lin MJ, Druso JE, Ramachandran S, et al. Liver-Type Glutaminase GLS2 Is a Druggable Metabolic Node in Luminal-Subtype Breast Cancer. Cell Rep. 2019;29(1):76-88 e7.
- de Visser KE, Joyce JA. The evolving tumor microenvironment: From cancer initiation to metastatic outgrowth. Cancer Cell. 2023;41(3):374–403.
- Vernieri C, Casola S, Foiani M, Pietrantonio F, de Braud F, Longo V. Targeting Cancer Metabolism: Dietary and Pharmacologic Interventions. Cancer Discov. 2016;6(12):1315–33.
- 12. Locasale JW, Goncalves MD, Di Tano M, Burgos-Barragan G. Diet and Cancer Metabolism. Cold Spring Harb Perspect Med. 2024;14:a041549.

- Locasale JW. Diet and Exercise in Cancer Metabolism. Cancer Discov. 2022;12(10):2249–57.
- Solanki S, Sanchez K, Ponnusamy V, Kota V, Bell HN, Cho CS, et al. Dysregulated Amino Acid Sensing Drives Colorectal Cancer Growth and Metabolic Reprogramming Leading to Chemoresistance. Gastroenterology. 2023;164(3):376-91 e13.
- Fontana L, Adelaiye RM, Rastelli AL, Miles KM, Ciamporcero E, Longo VD, et al. Dietary protein restriction inhibits tumor growth in human xenograft models. Oncotarget. 2013;4(12):2451–61.
- Lamming DW, Cummings NE, Rastelli AL, Gao F, Cava E, Bertozzi B, et al. Restriction of dietary protein decreases mTORC1 in tumors and somatic tissues of a tumor-bearing mouse xenograft model. Oncotarget. 2015;6(31):31233–40.
- Rubio-Patino C, Bossowski JP, De Donatis GM, Mondragon L, Villa E, Aira LE, et al. Low-Protein Diet Induces IRE1alpha-Dependent Anticancer Immunosurveillance. Cell Metab. 2018;27(4):828-42 e7.
- Upadhyayula PS, Higgins DM, Mela A, Banu M, Dovas A, Zandkarimi F, et al. Dietary restriction of cysteine and methionine sensitizes gliomas to ferroptosis and induces alterations in energetic metabolism. Nat Commun. 2023;14(1):1187.
- Maddocks ODK, Athineos D, Cheung EC, Lee P, Zhang T, van den Broek NJF, et al. Modulating the therapeutic response of tumours to dietary serine and glycine starvation. Nature. 2017;544(7650):372–6.
- Gao X, Sanderson SM, Dai Z, Reid MA, Cooper DE, Lu M, et al. Dietary methionine influences therapy in mouse cancer models and alters human metabolism. Nature. 2019;572(7769):397–401.
- Strekalova E, Malin D, Good DM, Cryns VL. Methionine Deprivation Induces a Targetable Vulnerability in Triple-Negative Breast Cancer Cells by Enhancing TRAIL Receptor-2 Expression. Clin Cancer Res. 2015;21(12):2780–91.
- Song M, Giovannucci E. Preventable Incidence and Mortality of Carcinoma Associated With Lifestyle Factors Among White Adults in the United States. JAMA Oncol. 2016;2(9):1154–61.
- Spencer CN, McQuade JL, Gopalakrishnan V, McCulloch JA, Vetizou M, Cogdill AP, et al. Dietary fiber and probiotics influence the gut microbiome and melanoma immunotherapy response. Science. 2021;374(6575):1632–40.
- Hoy AJ, Nagarajan SR, Butler LM. Tumour fatty acid metabolism in the context of therapy resistance and obesity. Nat Rev Cancer. 2021;21(12):753–66.
- Hopkins BD, Goncalves MD, Cantley LC. Obesity and Cancer Mechanisms: Cancer Metabolism. J Clin Oncol. 2016;34(35):4277–83.
- Peck B, Schulze A. Lipid Metabolism at the Nexus of Diet and Tumor Microenvironment. Trends Cancer. 2019;5(11):693–703.
- Hall KD, Guo J, Courville AB, Boring J, Brychta R, Chen KY, et al. Effect of a plant-based, low-fat diet versus an animal-based, ketogenic diet on ad libitum energy intake. Nat Med. 2021;27(2):344–53.
- Sanderson SM, Gao X, Dai Z, Locasale JW. Methionine metabolism in health and cancer: a nexus of diet and precision medicine. Nat Rev Cancer. 2019;19(11):625–37.
- 29. MacArthur MR, Mitchell SJ, Trevino-Villarreal JH, Grondin Y, Reynolds JS, Kip P, et al. Total protein, not amino acid composition, differs in plantbased versus omnivorous dietary patterns and determines metabolic health effects in mice. Cell Metab. 2021;33(9):1808-19 e2.
- Campbell TM, Campbell EK, Culakova E, Blanchard LM, Wixom N, Guido JJ, et al. A whole-food, plant-based randomized controlled trial in metastatic breast cancer: weight, cardiometabolic, and hormonal outcomes. Breast Cancer Res Treat. 2024;205(2):257–66.
- Campbell EK, Campbell TM, Culakova E, Blanchard L, Wixom N, Guido JJ, et al. A whole food, plant-based randomized controlled trial in metastatic breast cancer: feasibility, nutrient, and patient-reported outcomes. Breast Cancer Res Treat. 2024;206(2):273–83.
- 32. Melamud E, Vastag L, Rabinowitz JD. Metabolomic analysis and visualization engine for LC-MS data. Anal Chem. 2010;82(23):9818–26.
- Karlsen MC, Rogers G, Miki A, Lichtenstein AH, Folta SC, Economos CD, et al. Theoretical Food and Nutrient Composition of Whole-Food Plant-Based and Vegan Diets Compared to Current Dietary Recommendations. Nutrients. 2019;11(3):625.
- Ledikwe JH, Blanck HM, Kettel Khan L, Serdula MK, Seymour JD, Tohill BC, et al. Dietary energy density is associated with energy intake and weight status in US adults. Am J Clin Nutr. 2006;83(6):1362–8.

- Corbin KD, Carnero EA, Dirks B, Igudesman D, Yi F, Marcus A, et al. Hostdiet-gut microbiome interactions influence human energy balance: a randomized clinical trial. Nat Commun. 2023;14(1):3161.
- Obarzanek E, Sacks FM, Vollmer WM, Bray GA, Miller ER 3rd, Lin PH, et al. Effects on blood lipids of a blood pressure-lowering diet: the Dietary Approaches to Stop Hypertension (DASH) Trial. Am J Clin Nutr. 2001;74(1):80–9.
- Campbell TM, Campbell EK, Attia J, Ventura K, Mathews T, Chhabra KH, et al. The acute effects of a DASH diet and whole food, plant-based diet on insulin requirements and related cardiometabolic markers in individuals with insulin-treated type 2 diabetes. Diabetes Res Clin Pract. 2023;202:110814.
- Barnard ND, Cohen J, Jenkins DJ, Turner-McGrievy G, Gloede L, Jaster B, et al. A low-fat vegan diet improves glycemic control and cardiovascular risk factors in a randomized clinical trial in individuals with type 2 diabetes. Diabetes Care. 2006;29(8):1777–83.
- Barnard RJ, Lattimore L, Holly RG, Cherny S, Pritikin N. Response of noninsulin-dependent diabetic patients to an intensive program of diet and exercise. Diabetes Care. 1982;5(4):370–4.
- Simpson HC, Simpson RW, Lousley S, Carter RD, Geekie M, Hockaday TD, et al. A high carbohydrate leguminous fibre diet improves all aspects of diabetic control. Lancet. 1981;1(8210):1–5.
- Kazemi A, Speakman JR, Soltani S, Djafarian K. Effect of calorie restriction or protein intake on circulating levels of insulin like growth factor I in humans: A systematic review and meta-analysis. Clin Nutr. 2020;39(6):1705–16.
- Hidayat K, Chen JS, Wang TC, Liu YJ, Shi YJ, Su HW, et al. The Effects of Milk Supplementation on Bone Health Indices in Adults: A Meta-Analysis of Randomized Controlled Trials. Adv Nutr. 2022;13(4):1186–99.
- Harrison S, Lennon R, Holly J, Higgins JPT, Gardner M, Perks C, et al. Does milk intake promote prostate cancer initiation or progression via effects on insulin-like growth factors (IGFs)? A systematic review and metaanalysis. Cancer Causes Control. 2017;28(6):497–528.
- Linn T, Santosa B, Gronemeyer D, Aygen S, Scholz N, Busch M, et al. Effect of long-term dietary protein intake on glucose metabolism in humans. Diabetologia. 2000;43(10):1257–65.
- Tremblay F, Lavigne C, Jacques H, Marette A. Role of dietary proteins and amino acids in the pathogenesis of insulin resistance. Annu Rev Nutr. 2007;27:293–310.
- 46. Levine ME, Suarez JA, Brandhorst S, Balasubramanian P, Cheng CW, Madia F, et al. Low protein intake is associated with a major reduction in IGF-1, cancer, and overall mortality in the 65 and younger but not older population. Cell Metab. 2014;19(3):407–17.
- Fruman DA, Chiu H, Hopkins BD, Bagrodia S, Cantley LC, Abraham RT. The PI3K Pathway in Human Disease. Cell. 2017;170(4):605–35.
- Goncalves MD, Hopkins BD, Cantley LC. Phosphatidylinositol 3-Kinase, Growth Disorders, and Cancer. N Engl J Med. 2018;379(21):2052–62.
- Hopkins BD, Pauli C, Du X, Wang DG, Li X, Wu D, et al. Suppression of insulin feedback enhances the efficacy of PI3K inhibitors. Nature. 2018;560(7719):499–503.
- Cheung KL, Graves CR, Robertson JF. Tumour marker measurements in the diagnosis and monitoring of breast cancer. Cancer Treat Rev. 2000;26(2):91–102.
- Tajan M, Vousden KH. Dietary Approaches to Cancer Therapy. Cancer Cell. 2020;37(6):767–85.
- Niklison-Chirou MV, Erngren I, Engskog M, Haglöf J, Picard D, Remke M, et al. TAp73 is a marker of glutamine addiction in medulloblastoma. Genes Dev. 2017;31:1738.
- Green CL, Trautman ME, Chaiyakul K, Jain R, Alam YH, Babygirija R, et al. Dietary restriction of isoleucine increases healthspan and lifespan of genetically heterogeneous mice. Cell Metab. 2023;35(11):1976-95 e6.
- Taylor SR, Falcone JN, Cantley LC, Goncalves MD. Developing dietary interventions as therapy for cancer. Nat Rev Cancer. 2022;22(8):452–66.

#### **Publisher's Note**

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.